

Etabliert, zuverlässig  
und bewährt



## ImmunoCAP® ISAC

50 Jahre Erfahrung – konzentriert in einem Biochip

**ImmunoCAP ISAC** ist ein leistungsstarker und effizienter Immunoassay. Der Chip unterstützt Sie im klinischen Alltag, IgE-vermittelte Allergien zu diagnostizieren, zu verstehen und zu behandeln.

**Hohe Zuverlässigkeit** in einer globalen Validierungsstudie gezeigt.<sup>206</sup>

**Breiter Nutzen** in über 220 Veröffentlichungen beschrieben.<sup>1-226</sup>

**Der klinische Nutzen des ImmunoCAP ISAC ist für zahlreiche allergische Erkrankungen nachgewiesen.**<sup>149</sup>

- Atopische Dermatitis<sup>41, 65, 78, 121, 145, 162</sup>
- Asthma und Rhinitis<sup>22, 47, 103, 138, 150, 151, 191</sup>
- Idiopathische Anaphylaxie<sup>88</sup>
- Kofaktor-abhängige Nahrungsmittel-Allergie<sup>35, 147, 163</sup>
- Indikation für eine spezifische Immuntherapie<sup>12, 108, 173, 180, 205, 223</sup>
- Eosinophile Ösophagitis<sup>10, 61, 190, 209</sup>
- Risikoeinschätzung und -bewertung<sup>11, 87, 139, 214</sup>

**Wünschen Sie weitere Informationen?**  
Sprechen Sie uns gerne an.

**ThermoFisher**  
SCIENTIFIC

# Referenzen, 2002–2017

(Basierend auf den Suchergebnissen unter [www.ncbi.nlm.nih.gov/pubmed](http://www.ncbi.nlm.nih.gov/pubmed))

1. Ackerbauer, D., et al., *Component-resolved IgE profiles in Austrian patients with a convincing history of peanut allergy*. Int Arch Allergy Immunol, 2015. 166(1): p. 13-24.
2. Agabriel, C., et al., *Ara h 2 and Ara h 6 sensitization predicts peanut allergy in Mediterranean pediatric patients*. Pediatr Allergy Immunol, 2014. 25(7): p. 662-7.
3. Ahlgrim, C., et al., *Comparison of Molecular Multiplex and Singleplex Analysis of IgE to Grass Pollen Allergens in Untreated German Grass Pollen-Allergic Patients*. J Investig Allergol Clin Immunol, 2015. 25(3): p. 190-5.
4. Alessandri, C., et al., *Ovomucoid (Gal d 1) specific IgE detected by microarray system predict tolerability to boiled hen's egg and an increased risk to progress to multiple environmental allergen sensitisation*. Clin Exp Allergy, 2012. 42(3): p. 441-50.
5. Alvarado, M.I., et al., *Profilin as a severe food allergen in allergic patients overexposed to grass pollen*. Allergy, 2014. 69(12): p. 1610-6.
6. Amo, A., et al., *Gal d 6 is the second allergen characterized from egg yolk*. J Agric Food Chem, 2010. 58(12): p. 7453-7.
7. Araujo, L.M., N.A. Rosario, and A. Mari, *Molecular-based diagnosis of respiratory allergic diseases in children from Curitiba, a city in Southern Brazil*. Allergol Immunopathol (Madr), 2016. 44(1): p. 18-22.
8. Ariano, R., et al., *Occupational asthma associated to the exposure to limonium tataricum flowers*. Eur Ann Allergy Clin Immunol, 2013. 45(3): p. 84-9.
9. Armentia, A., et al., *Value of microarray allergen assay in the management of eosinophilic oesophagitis*. Allergol Immunopathol (Madr), 2015. 43(1): p. 73-80.
10. Armentia, A., et al., *Anaphylaxis caused by hidden soybean allergens in pillows*. J Allergy Clin Immunol, 2013. 131(1): p. 228-30.
11. Asarnoj, A., et al., *Sensitization to cat and dog allergen molecules in childhood and prediction of symptoms of cat and dog allergy in adolescence: A BAMSE/MeDALL study*. J Allergy Clin Immunol, 2016. 137(3): p. 813-21 e7.
12. Asero, R., *Component-resolved diagnosis-assisted prescription of allergen-specific immunotherapy: a practical guide*. Eur Ann Allergy Clin Immunol, 2012. 44(5): p. 183-7.
13. Azofra, J., et al., *Lipid Transfer Protein Syndrome in a Non-Mediterranean Area*. Int Arch Allergy Immunol, 2016. 169(3): p. 181-8.
14. Barićić, T.V. and S. Dodig, *Birch pollen-associated peanut allergies in children*. Aerobiologia, 2013. 29(1): p. 85-93.
15. Barićić, T.V. and S. Dodig, *Genuine and Cross-Reactive Peanut Components in Birch-Sensitive Preschool Children*. Laboratory Medicine, 2013. 44(3): p. 210-214.
16. Baron-Bodo, V., et al., *Absence of IgE neosensitization in house dust mite allergic patients following sublingual immunotherapy*. Clin Exp Allergy, 2012. 42(10): p. 1510-8.
17. Batard, T., et al., *Patterns of IgE sensitization in house dust mite-allergic patients: implications for allergen immunotherapy*. Allergy, 2016. 71(2): p. 220-9.
18. Becker, S., et al., *Non-allergic rhinitis with eosinophilia syndrome is not associated with local production of specific IgE in nasal mucosa*. Eur Arch Otorhinolaryngol, 2016. 273(6): p. 1469-75.
19. Becker, S., et al., *Real-Life Study for the Diagnosis of House Dust Mite Allergy - The Value of Recombinant Allergen-Based IgE Serology*. Int Arch Allergy Immunol, 2016. 170(2): p. 132-7.
20. Berneder, M., et al., *Allergen chip diagnosis for soy-allergic patients: Gly m 4 as a marker for severe food-allergic reactions to soy*. Int Arch Allergy Immunol, 2013. 161(3): p. 229-33.
21. Bienvenu, J., P. Rouzaire, and F. Bienvenu, *Les allergènes moléculaires : évolution ou révolution dans le diagnostic de l'allergie*. Revue Française d'Allergologie, 2011. 51(3): p. 186-191.
22. Bjerg, A., et al., *A population-based study of animal component sensitization, asthma, and rhinitis in schoolchildren*. Pediatr Allergy Immunol, 2015. 26(6): p. 557-63.
23. Bokanovic, D., et al., *Determination of slgE to rPhl p 1 is sufficient to diagnose grass pollen allergy*. Allergy, 2013. 68(11): p. 1403-9.
24. Bonini, M., et al., *Microarray evaluation of specific IgE to allergen components in elite athletes*. Allergy, 2012. 67(12): p. 1557-64.
25. Borres, M.P., M. Ebisawa, and P.A. Eigenmann, *Use of allergen components begins a new era in pediatric allergology*. Pediatr Allergy Immunol, 2011. 22(5): p. 454-61.
26. Bousquet, J., et al., *Are allergic multimorbidities and IgE polysensitization associated with the persistence or re-occurrence of foetal type 2 signalling? The MeDALL hypothesis*. Allergy, 2015. 70(9): p. 1062-78.
27. Brans, R., et al., *Microarray-based detection of specific IgE against recombinant omega-5-gliadin in suspected wheat-dependent exercise-induced anaphylaxis*. Eur J Dermatol, 2012. 22(3): p. 358-62.
28. Bronnert, M., et al., *Component-resolved diagnosis with commercially available D. pteronyssinus Der p 1, Der p 2 and Der p 10: relevant markers for house dust mite allergy*. Clin Exp Allergy, 2012. 42(9): p. 1406-15.
29. Burney, P.G., et al., *The prevalence and distribution of food sensitization in European adults*. Allergy, 2014. 69(3): p. 365-71.
30. Cabauatan, C.R., et al., *Allergen microarray detects high prevalence of asymptomatic IgE sensitizations to tropical pollen-derived carbohydrates*. J Allergy Clin Immunol, 2014. 133(3): p. 910-4 e5.
31. Cabrera-Freitag, P., et al., *Can component-based microarray replace fluorescent enzimoimmunoassay in the diagnosis of grass and cypress pollen allergy?* Clin Exp Allergy, 2011. 41(10): p. 1440-6.
32. Cabrera-Freitag, P., et al., *A study of the variability of the in vitro component-based microarray ISAC CDR 103 technique*. J Investig Allergol Clin Immunol, 2011. 21(5): p. 414-5.
33. Caminati, M., et al., *Component resolved diagnosis (CRD): how much is it presently used by Italian allergists?* Eur Ann Allergy Clin Immunol, 2014. 46(3): p. 106-8.
34. Canonica, G.W., et al., *A WAO - ARIA - GA(2)LEN consensus document on molecular-based allergy diagnostics*. World Allergy Organ J, 2013. 6(1): p. 17.

35. Cardona, V., et al., *Co-factor-enhanced food allergy*. Allergy, 2012. 67(10): p. 1316-8.
36. Caubet, J.C. and H.A. Sampson, *Beyond skin testing: state of the art and new horizons in food allergy diagnostic testing*. Immunol Allergy Clin North Am, 2012. 32(1): p. 97-109.
37. Chambel, M., et al., *Polissensibilização a pólenes analisada e reinterpretada à luz do método ImmunoCAP ISAC®*. Revista Portuguesa de Imunoalergologia, 2012. 20: p. 211-219.
38. Chawes, B.L., *Low-grade disease activity in early life precedes childhood asthma and allergy*. Dan Med J, 2016. 63(8).
39. Chelminska, M., et al., *Differentiating of cross-reactions in patients with latex allergy with the use of ISAC test*. Postepy Dermatol Alergol, 2016. 33(2): p. 120-7.
40. Chen, Q., et al., *Allergic sensitization patterns identified through latent class analysis among children with and without asthma*. Ann Allergy Asthma Immunol, 2016. 116(3): p. 212-8.
41. Choi, J.S., J.Y. Roh, and J.R. Lee, *Clinical availability of component-resolved diagnosis using microarray technology in atopic dermatitis*. Ann Dermatol, 2014. 26(4): p. 437-46.
42. Ciprandi, G., et al., *Walnut anaphylaxis: the usefulness of molecular-based allergy diagnostics*. Immunol Lett, 2014. 161(1): p. 138-9.
43. Ciprandi, G., et al., *Peanut anaphylaxis: the usefulness of molecular-based allergy diagnostics*. Allergy, 2015. 70(1): p. 129-30.
44. Couto, M., et al., *Anaphylaxis to sunflower seed*. Rev Port Imunoalergol, 2012. 20(2): p. 139-143.
45. Cretich, M., F. Damin, and M. Chiari, *Protein microarray technology: how far off is routine diagnostics?* Analyst, 2014. 139(3): p. 528-42.
46. Curin, M., et al., *Microarrayed dog, cat, and horse allergens show weak correlation between allergen-specific IgE and IgG responses*. J Allergy Clin Immunol, 2014. 133(3): p. 918-21 e6.
47. Custovic, A., et al., *Evolution pathways of IgE responses to grass and mite allergens throughout childhood*. J Allergy Clin Immunol, 2015. 136(6): p. 1645-52 e1-8.
48. da Silva, D.M., et al., *Cross-reactive LTP sensitization in food-dependent exercise-induced urticaria/anaphylaxis: a pilot study of a component-resolved and in vitro depletion approach*. Clin Transl Allergy, 2016. 6: p. 46.
49. D'Amelio, C.M., et al., *Is the performance of ImmunoCAP ISAC 112 sufficient to diagnose peach and apple allergies?* Ann Allergy Asthma Immunol, 2016. 116(2): p. 162-3.
50. Dang, T.D., C.E. Mills, and K.J. Allen, *Determination of the clinical egg allergy phenotypes using component-resolved diagnostics*. Pediatr Allergy Immunol, 2014. 25(7): p. 639-43.
51. De Knop, K.J., et al., *Age-related sensitization profiles for hazelnut (*Corylus avellana*) in a birch-endemic region*. Pediatr Allergy Immunol, 2011. 22(1 Pt 2): p. e139-49.
52. De Linares, C., et al., *Optimization of the measurement of outdoor airborne allergens using a protein microarrays platform*. Aerobiologia, 2014. 30(3): p. 217-227.
53. De Swert, L.F., et al., *Secondary soy allergy in children with birch pollen allergy may cause both chronic and acute symptoms*. Pediatr Allergy Immunol, 2012. 23(2): p. 117-23.
54. Deinhofer, K., et al., *Microarrayed allergens for IgE profiling*. Methods, 2004. 32(3): p. 249-54.
55. Dollner, R., et al., *Non-allergic, mastocytosis-associated rhinitis*. Clin Exp Allergy, 2013. 43(4): p. 406-12.
56. D'Urbano, L.E., et al., *Performance of a component-based allergen-microarray in the diagnosis of cow's milk and hen's egg allergy*. Clin Exp Allergy, 2010. 40(10): p. 1561-70.
57. Ebo, D.G., et al., *Sensitization profiles in birch pollen-allergic patients with and without oral allergy syndrome to apple: lessons from multiplexed component-resolved allergy diagnosis*. Clin Exp Allergy, 2010. 40(2): p. 339-47.
58. Ebo, D.G., et al., *Component-resolved diagnosis from latex allergy by microarray*. Clin Exp Allergy, 2010. 40(2): p. 348-58.
59. Ebo, D.G., et al., *New food allergies in a European non-Mediterranean region: is Cannabis sativa to blame?* Int Arch Allergy Immunol, 2013. 161(3): p. 220-8.
60. Eigenmann, P.A., *Component-resolved diagnosis in food allergy, are micro-array assays helpful to the clinician?* Allergy, 2008. 63(11): p. 1519-20.
61. Erwin, E.A., et al., *Serum IgE measurement and detection of food allergy in pediatric patients with eosinophilic esophagitis*. Ann Allergy Asthma Immunol, 2010. 104(6): p. 496-502.
62. Erwin, E.A., et al., *IgE Antibody Detection and Component Analysis in Patients with Eosinophilic Esophagitis*. J Allergy Clin Immunol Pract, 2015. 3(6): p. 896-904 e3.
63. Ferrer, M., et al., *Molecular diagnosis in allergology: application of the microarray technique*. J Investig Allergol Clin Immunol, 2009. 19 Suppl 1: p. 19-24.
64. Foong, R.X., et al., *Pilot study: assessing the clinical diagnosis of allergy in atopic children using a microarray assay in addition to skin prick testing and serum specific IgE*. Clin Mol Allergy, 2016. 14: p. 8.
65. Fung, I., J.S. Kim, and J.M. Spergel, *Relating microarray component testing and reported food allergy and food-triggered atopic dermatitis: a real-world analysis*. Ann Allergy Asthma Immunol, 2013. 110(3): p. 173-177 e1.
66. Gadisseur, R., J.P. Chapelle, and E. Cavalier, *A new tool in the field of in-vitro diagnosis of allergy: preliminary results in the comparison of ImmunoCAP(c) 250 with the ImmunoCAP(c) ISAC*. Clin Chem Lab Med, 2011. 49(2): p. 277-80.
67. Garcia, B.E., et al., *Is the ISAC 112 Microarray Useful in the Diagnosis of Pollinosis in Spain?* J Investig Allergol Clin Immunol, 2016. 26(2): p. 92-9.
68. Garcia-Ara, C., et al., *Sensitization to microarrayed species-specific plant components precedes that of cross-reacting allergens*. Pediatr Allergy Immunol, 2017. 28(3): p. 288-291.
69. Garcia-Blanca, A., et al., *Influence of age on IgE response in peanut-allergic children and adolescents from the Mediterranean area*. Pediatr Allergy Immunol, 2015. 26(6): p. 497-502.
70. Gassner, M., R. Gehrig, and P. Schmid-Grendelmeier, *Hay fever as a Christmas gift*. N Engl J Med, 2013. 368(4): p. 393-4.
71. Geroldinger-Simic, M., et al., *Oral exposure to Mal d 1 affects the immune response in patients with birch pollen allergy*. J Allergy Clin Immunol, 2013. 131(1): p. 94-102.
72. Giuffrida, M.G., et al., *Shrimp allergy beyond Tropomyosin in Italy: clinical relevance of Arginine Kinase, Sarcoplasmic calcium binding protein and Hemocyanin*. Eur Ann Allergy Clin Immunol, 2014. 46(5): p. 172-7.
73. Goikoetxea, M.J., et al., *The importance of in vitro component-resolved diagnosis in paediatric patients*. Allergol Immunopathol (Madr), 2010. 38(1): p. 37-40.

74. Goikoetxea, M.J., et al., *Is Microarray Analysis Really Useful and Sufficient to Diagnose Nut Allergy in the Mediterranean Area?* J Investig Allergol Clin Immunol, 2016. 26(1): p. 31-9.
75. Gray, C.L., M.E. Levin, and G. du Toit, *Ethnic differences in peanut allergy patterns in South African children with atopic dermatitis.* Pediatr Allergy Immunol, 2015. 26(8): p. 721-30.
76. Gray, C.L., M.E. Levin, and G. du Toit, *Egg sensitization, allergy and component patterns in African children with atopic dermatitis.* Pediatr Allergy Immunol, 2016. 27(7): p. 709-715.
77. Gray, C.L., M.E. Levin, and G. Du Toit, *Which test is best for diagnosing peanut allergy in South African children with atopic dermatitis?* S Afr Med J, 2016. 106(2): p. 214-20.
78. Gray, C.L., et al., *Food allergy in South African children with atopic dermatitis.* Pediatr Allergy Immunol, 2014. 25(6): p. 572-9.
79. Griffiths, R.L.M., et al., *Comparison of the Performance of Skin Prick, ImmunoCAP, and ISAC Tests in the Diagnosis of Patients with Allergy.* Int Arch Allergy Immunol, 2017. 172(4): p. 215-223.
80. Hamilton, R.G., *Microarray Technology Applied to Human Allergic Disease.* Microarrays (Basel), 2017. 6(1).
81. Hamilton, R.G. and J. Kleine-Tebbe, *Molecular Allergy Diagnostics: Analytical Features That Support Clinical Decisions.* Curr Allergy Asthma Rep, 2015. 15(9): p. 57.
82. Hantusch, B., et al., *Affinity determinations of purified IgE and IgG antibodies against the major pollen allergens Phl p 5a and Bet v 1a: discrepancy between IgE and IgG binding strength.* Immunol Lett, 2005. 97(1): p. 81-9.
83. Harwanegg, C. and R. Hiller, *Protein microarrays in diagnosing IgE-mediated diseases: spotting allergy at the molecular level.* Expert Rev Mol Diagn, 2004. 4(4): p. 539-48.
84. Harwanegg, C. and R. Hiller, *Protein microarrays for the diagnosis of allergic diseases: state-of-the-art and future development.* Clin Chem Lab Med, 2005. 43(12): p. 1321-6.
85. Harwanegg, C., et al., *Microarrayed recombinant allergens for diagnosis of allergy.* Clin Exp Allergy, 2003. 33(1): p. 7-13.
86. Hatzler, L., et al., *Parental hay fever reinforces IgE to pollen as pre-clinical biomarker of hay fever in childhood.* Pediatr Allergy Immunol, 2014. 25(4): p. 366-73.
87. Hatzler, L., et al., *Molecular spreading and predictive value of preclinical IgE response to *Phleum pratense* in children with hay fever.* J Allergy Clin Immunol, 2012. 130(4): p. 894-901 e5.
88. Heaps, A., et al., *The utility of the ISAC allergen array in the investigation of idiopathic anaphylaxis.* Clin Exp Immunol, 2014. 177(2): p. 483-90.
89. Hedlin, G., J. Konradsen, and A. Bush, *An update on paediatric asthma.* European Respiratory Review, 2012. 21(125): p. 175-185.
90. Hiller, R., et al., *Microarrayed allergen molecules: diagnostic gatekeepers for allergy treatment.* FASEB J, 2002. 16(3): p. 414-6.
91. Hochwallner, H., et al., *Transmission of allergen-specific IgG and IgE from maternal blood into breast milk visualized with microarray technology.* J Allergy Clin Immunol, 2014. 134(5): p. 1213-5.
92. Holzweber, F., et al., *Inhibition of IgE binding to cross-reactive carbohydrate determinants enhances diagnostic selectivity.* Allergy, 2013. 68(10): p. 1269-77.
93. Hong, X., et al., *IgE, but not IgG4, antibodies to Ara h 2 distinguish peanut allergy from asymptomatic peanut sensitization.* Allergy, 2012. 67(12): p. 1538-46.
94. Huss-Marp, J., et al., *Comparison of molecular and extract-based allergy diagnostics with multiplex and singleplex analysis.* Allergo J Int, 2015. 24: p. 46-53.
95. Incorvaia, C., et al., *A Pitfall to Avoid When Using an Allergen Microarray: The Incidental Detection of IgE to Unexpected Allergens.* J Allergy Clin Immunol Pract, 2015. 3(6): p. 879-82.
96. Jahn-Schmid, B., et al., *Allergen microarray: comparison of microarray using recombinant allergens with conventional diagnostic methods to detect allergen-specific serum immunoglobulin E.* Clin Exp Allergy, 2003. 33(10): p. 1443-9.
97. Javaloyes, G., et al., *Performance of different in vitro techniques in the molecular diagnosis of peanut allergy.* J Investig Allergol Clin Immunol, 2012. 22(7): p. 508-13.
98. Keet, C.A., et al., *Evaluation of Ara h2 IgE thresholds in the diagnosis of peanut allergy in a clinical population.* J Allergy Clin Immunol Pract, 2013. 1(1): p. 101-3.
99. Klemans, R.J., et al., *The diagnostic accuracy of specific IgE to Ara h 6 in adults is as good as Ara h 2.* Allergy, 2014. 69(8): p. 1112-4.
100. Klemans, R.J., et al., *IgE binding to peanut components by four different techniques: Ara h 2 is the most relevant in peanut allergic children and adults.* Clin Exp Allergy, 2013. 43(8): p. 967-74.
101. Klimek, L., et al., *[Microarray technique for component resolved diagnosis (CRD) in type-I allergies. An innovative technology at the border between research tool and routine diagnostics].* HNO, 2011. 59(10): p. 988-93.
102. Knol, E.F. and A.C. Knulst, *Application of multiplexed immunoglobulin E determination on a chip in component-resolved diagnostics in allergy.* Clin Exp Allergy, 2010. 40(2): p. 190-2.
103. Konradsen, J.R., et al., *Severe childhood asthma and allergy to furry animals: refined assessment using molecular-based allergy diagnostics.* Pediatr Allergy Immunol, 2014. 25(2): p. 187-92.
104. Krause, S., et al., *Lipid transfer protein (Ara h 9) as a new peanut allergen relevant for a Mediterranean allergic population.* J Allergy Clin Immunol, 2009. 124(4): p. 771-8 e5.
105. Kukkonen, A.K., et al., *Turnip rapeseed does not worsen atopic eczema in most sensitized children.* Pediatr Allergy Immunol, 2014. 25(6): p. 613-5.
106. Kukkonen, A.K., et al., *Ara h 2 and Ara 6 are the best predictors of severe peanut allergy: a double-blind placebo-controlled study.* Allergy, 2015. 70(10): p. 1239-45.
107. Leonardi, A., et al., *Microarray-based IgE detection in tears of patients with vernal keratoconjunctivitis.* Pediatr Allergy Immunol, 2015. 26(7): p. 641-5.
108. Letran, A., M. Espinazo, and F. Moreno, *Measurement of IgE to pollen allergen components is helpful in selecting patients for immunotherapy.* Ann Allergy Asthma Immunol, 2013. 111(4): p. 295-7.
109. Liccardi, G., et al., *What could be the role of molecular-based allergy diagnostics in detecting the risk of developing allergic sensitization to furry animals?* Eur Ann Allergy Clin Immunol, 2015. 47(5): p. 163-7.
110. Liccardi, G., et al., *Sensitization to Common and Uncommon Pets or Other Furry Animals: Which May Be Common Mechanisms?* Transl Med UniSa, 2016. 14: p. 9-14.
111. Lidholm, J., et al., *Component-resolved diagnostics in food allergy.* Curr Opin Allergy Clin Immunol, 2006. 6(3): p. 234-40.
112. Lizaso, M.T., et al., *Comparison of conventional and component-resolved diagnostics by two different methods (Advia-Centaur/Microarray-ISAC) in pollen allergy.* Ann Allergy Asthma Immunol, 2011. 107(1): p. 35-41

113. Lomidze, N. and M. Gotua, *Patterns of Sensitization by Food and Inhalant Components in Georgian Patients*. Georgian Med News, 2016(259): p. 23-26.
114. Lopes de Oliveira, L.C., et al., *The value of specific IgE to peanut and its component Ara h 2 in the diagnosis of peanut allergy*. J Allergy Clin Immunol Pract, 2013. 1(4): p. 394-8.
115. Lucas, J.M., *Microarrays: Molecular allergology and nanotechnology for personalised medicine (II)*. Allergol Immunopathol (Madr), 2010. 38(4): p. 217-23.
116. Lucas, J.M., *Microarrays: molecular allergology and nanotechnology for personalised medicine (I)*. Allergol Immunopathol (Madr), 2010. 38(3): p. 153-61.
117. Lupinek, C., et al., *Analysis of serum IgE reactivity profiles with microarrayed allergens indicates absence of de novo IgE sensitizations in adults*. J Allergy Clin Immunol, 2012. 130(6): p. 1418-20 e4.
118. Lupinek, C., et al., *Advances in allergen-microarray technology for diagnosis and monitoring of allergy: the MeDALL allergen-chip*. Methods, 2014. 66(1): p. 106-19.
119. Macchia, D., et al., *Guidelines for the use and interpretation of diagnostic methods in adult food allergy*. Clin Mol Allergy, 2015. 13: p. 27.
120. Mari, A., et al., *Microarrayed allergen molecules for the diagnosis of allergic diseases*. Curr Allergy Asthma Rep, 2010. 10(5): p. 357-64.
121. Mari, A., E. Scala, and C. Alessandri, *The IgE-microarray testing in atopic dermatitis: a suitable modern tool for the immunological and clinical phenotyping of the disease*. Curr Opin Allergy Clin Immunol, 2011. 11(5): p. 438-44.
122. Martinez-Aranguren, R., et al., *Is the determination of specific IgE against components using ISAC 112 a reproducible technique?* PLoS One, 2014. 9(2): p. e88394.
123. Martinez-Aranguren, R., et al., *Clinical Performance of Commercial ISAC 112 Allergen Microarray Versus Noncommercial RIRAAF Platform for the Diagnosis of Plant Food and Olive Pollen Allergies*. J Investig Allergol Clin Immunol, 2016. 26(3): p. 185-7.
124. Mauro, M., et al., *Birch-apple syndrome treated with birch pollen immunotherapy*. Int Arch Allergy Immunol, 2011. 156(4): p. 416-22.
125. Meglio, P., et al., *Oral food desensitization in children with IgE-mediated hen's egg allergy: a new protocol with raw hen's egg*. Pediatr Allergy Immunol, 2013. 24(1): p. 75-83.
126. Melioli, G., et al., *The ImmunoCAP ISAC molecular allergology approach in adult multi-sensitized Italian patients with respiratory symptoms*. Clin Biochem, 2011. 44(12): p. 1005-11.
127. Melioli, G. and G.W. Canonica, *Molecular allergy diagnosis: we need to become more knowledgeable*. Ann Allergy Asthma Immunol, 2012. 108(6): p. 387.
128. Melioli, G., et al., *The added value of allergen microarray technique to the management of poly-sensitized allergic patients*. Curr Opin Allergy Clin Immunol, 2012. 12(4): p. 434-9.
129. Melioli, G., et al., *The IgE repertoire in children and adolescents resolved at component level: a cross-sectional study*. Pediatr Allergy Immunol, 2012. 23(5): p. 433-40.
130. Melioli, G., et al., *Allergenius, an expert system for the interpretation of allergen microarray results*. World Allergy Organ J, 2014. 7(1): p. 15.
131. Mercier, V., *Apports et limites du diagnostic moléculaire dans la prise en charge des allergies alimentaires*. Revue Française d'Allergologie, 2012. 52: p. S19-S26.
132. Metz-Favre, C., et al., *Molecular allergology in practice: an unusual case of LTP allergy*. Eur Ann Allergy Clin Immunol, 2011. 43(6): p. 193-5.
133. Mohamad Yadzir, Z.H., et al., *Improved Diagnosis of the Polysensitized Allergic Rhinitis Patients Using Component Resolved Diagnosis Method*. Iran J Allergy Asthma Immunol, 2016. 15(2): p. 156-60.
134. Mohamad Yadzir, Z.H., et al., *Component-resolved diagnosis (CRD): Is it worth it? frequency and differentiation in rhinitis patients with mite reactivity*. Iran J Allergy Asthma Immunol, 2014. 13(4): p. 240-6.
135. Moneret-Vautrin, D.A., et al., *Component resolved diagnosis by the ISAC allergen microarray*. Revue Francaise d'Allergologie, 2011. 51(2): p. 73-83.
136. Nettis, E., et al., *Molecular diagnosis and the Italian Board for ISAC*. Eur Ann Allergy Clin Immunol, 2014. 46(2): p. 68-73.
137. Nolte, M., et al., *Timothy specific IgE levels are associated with efficacy and safety of timothy grass sublingual immunotherapy tablet*. Ann Allergy Asthma Immunol, 2015. 115(6): p. 509-515 e2.
138. Nordlund, B., et al., *IgE antibodies to animal-derived lipocalin, kallikrein and secretoglobin are markers of bronchial inflammation in severe childhood asthma*. Allergy, 2012. 67(5): p. 661-9.
139. Onell, A., L. Hjalle, and M.P. Borres, *Exploring the temporal development of childhood IgE profiles to allergen components*. Clin Transl Allergy, 2012. 2(1): p. 24.
140. Onell, A., et al., *Allergy testing in children with persistent asthma: comparison of four diagnostic methods*. Allergy, 2017. 72(4): p. 590-597.
141. Ott, H., et al., *Clinical usefulness of microarray-based IgE detection in children with suspected food allergy*. Allergy, 2008. 63(11): p. 1521-8.
142. Ott, H., et al., *Allergen microarrays: a novel tool for high-resolution IgE profiling in adults with atopic dermatitis*. Eur J Dermatol, 2010. 20(1): p. 54-61.
143. Ott, H., et al., *Microarrays of recombinant Hevea brasiliensis proteins: a novel tool for the component-resolved diagnosis of natural rubber latex allergy*. J Investig Allergol Clin Immunol, 2010. 20(2): p. 129-38.
144. Ott, H., et al., *Microarray-based IgE detection in capillary blood samples of patients with atopy*. Allergy, 2006. 61(9): p. 1146-7.
145. Ott, H., et al., *Molecular microarray analysis reveals allergen- and exotoxin-specific IgE repertoires in children with atopic dermatitis*. J Eur Acad Dermatol Venereol, 2014. 28(1): p. 100-7.
146. Panzner, P., et al., *A comprehensive analysis of middle-European molecular sensitization profiles to pollen allergens*. Int Arch Allergy Immunol, 2014. 164(1): p. 74-82.
147. Pascal, M., et al., *Lipid transfer protein syndrome: clinical pattern, cofactor effect and profile of molecular sensitization to plant-foods and pollens*. Clin Exp Allergy, 2012. 42(10): p. 1529-39.
148. Passalacqua, G., et al., *The additional values of microarray allergen assay in the management of polysensitized patients with respiratory allergy*. Allergy, 2013. 68(8): p. 1029-33.
149. Patelis, A., et al., *Multiplex component-based allergen microarray in recent clinical studies*. Clin Exp Allergy, 2016. 46(8): p. 1022-32.
150. Patelis, A., et al., *Allergen extract vs. component sensitization and airway inflammation, responsiveness and new-onset respiratory disease*. Clin Exp Allergy, 2016. 46(5): p. 730-40.
151. Patelis, A., et al., *Population-based study of multiplexed IgE sensitization in relation to asthma, exhaled nitric oxide, and bronchial responsiveness*. J Allergy Clin Immunol, 2012. 130(2): p. 397-402 e2.

152. Pauli, G. and T. Chivato, *Allergologie moléculaire en pratique : à propos d'un patient polysensibilisé présentant plusieurs allergies alimentaires sévères*. Revue Française d'Allergologie, 2010. 50(6): p. 513-515.
153. Pedrosa, M., et al., *Utility of specific IgE to Ara h 6 in peanut allergy diagnosis*. Ann Allergy Asthma Immunol, 2015. 115(2): p. 108-12.
154. Pedrosa, M., et al., *Peanut seed storage proteins are responsible for clinical reactivity in Spanish peanut-allergic children*. Pediatr Allergy Immunol, 2012. 23(7): p. 654-9.
155. Pfiffner, P., et al., *Cross-reactions vs co-sensitization evaluated by in silico motifs and in vitro IgE microarray testing*. Allergy, 2012. 67(2): p. 210-6.
156. Prosperi, M.C., et al., *Challenges in interpreting allergen microarrays in relation to clinical symptoms: a machine learning approach*. Pediatr Allergy Immunol, 2014. 25(1): p. 71-9.
157. Quirce, S., *IgE antibodies in occupational asthma: are they causative or an associated phenomenon?* Curr Opin Allergy Clin Immunol, 2014. 14(2): p. 100-5.
158. Rashid, R.S., et al., *Pollen-food syndrome is related to Bet v 1/PR-10 protein sensitisation, but not all patients have spring rhinitis*. Allergy, 2011. 66(10): p. 1391-2.
159. Reisacher, W.R., *Detecting local immunoglobulin E from mucosal brush biopsy of the inferior turbinates using microarray analysis*. Int Forum Allergy Rhinol, 2013. 3(5): p. 399-403.
160. Rentzos, G., et al., *Intestinal allergic inflammation in birch pollen allergic patients in relation to pollen season, IgE sensitization profile and gastrointestinal symptoms*. Clin Transl Allergy, 2014. 4: p. 19.
161. Resch, Y., et al., *Different IgE recognition of mite allergen components in asthmatic and nonasthmatic children*. J Allergy Clin Immunol, 2015. 136(4): p. 1083-91.
162. Rockmann, H., et al., *Food allergen sensitization pattern in adults in relation to severity of atopic dermatitis*. Clin Transl Allergy, 2014. 4(1): p. 9.
163. Romano, A., et al., *Lipid transfer proteins: the most frequent sensitizer in Italian subjects with food-dependent exercise-induced anaphylaxis*. Clin Exp Allergy, 2012. 42(11): p. 1643-53.
164. Rossi, R.E., et al., *Sensitization profiles in polysensitized patients from a restricted geographical area: further lessons from multiplexed component resolved diagnosis*. Eur Ann Allergy Clin Immunol, 2011. 43(6): p. 171-5.
165. Ruiz-Garcia, M., et al., *Profilin: a relevant aeroallergen?* J Allergy Clin Immunol, 2011. 128(2): p. 416-8.
166. Salcedo, G. and A. Diaz-Perales, *Component-resolved diagnosis of allergy: more is better?* Clin Exp Allergy, 2010. 40(6): p. 836-8.
167. Sano, A., et al., *Sensitization profiles of a case of pollen-food allergy syndrome*. Allergol Int, 2011. 60(1): p. 97-101.
168. Santos, N., et al., *Anaphylaxis to apple*. Rev Port Imunoalergol, 2011. 19(4): p. 223-227.
169. Santosa, A., et al., *Evaluation of the applicability of the Immuno-solid-phase allergen chip (ISAC) assay in atopic patients in Singapore*. Clin Transl Allergy, 2015. 5: p. 9.
170. Sanz, M.L., A.B. Blazquez, and B.E. Garcia, *Microarray of allergenic component-based diagnosis in food allergy*. Curr Opin Allergy Clin Immunol, 2011. 11(3): p. 204-9.
171. Sastre, J., *Molecular diagnosis in allergy*. Clin Exp Allergy, 2010. 40(10): p. 1442-60.
172. Sastre, J., *Molecular diagnosis and immunotherapy*. Curr Opin Allergy Clin Immunol, 2013. 13(6): p. 646-50.
173. Sastre, J., et al., *How molecular diagnosis can change allergen-specific immunotherapy prescription in a complex pollen area*. Allergy, 2012. 67(5): p. 709-11.
174. Sastre, J., et al., *Adverse reactions to immunotherapy are associated with different patterns of sensitization to grass allergens*. Allergy, 2015. 70(5): p. 598-600.
175. Scala, E., et al., *Molecular Recognition Profiles and Clinical Patterns of PR-10 Sensitization in a Birch-Free Mediterranean Area*. International Archives of Allergy and Immunology, 2017: p. 138-146.
176. Scala, E., et al., *Specific IgE toward allergenic molecules is a new prognostic marker in patients with Sezary syndrome*. Int Arch Allergy Immunol, 2012. 157(2): p. 159-67.
177. Scala, E., et al., *Cross-sectional survey on immunoglobulin E reactivity in 23,077 subjects using an allergenic molecule-based microarray detection system*. Clin Exp Allergy, 2010. 40(6): p. 911-21.
178. Scala, E., et al., *IgE recognition patterns of profilin, PR-10, and tropomyosin panallergens tested in 3,113 allergic patients by allergen microarray-based technology*. PLoS One, 2011. 6(9): p. e24912.
179. Scala, E., et al., *Lipid transfer protein sensitization: reactivity profiles and clinical risk assessment in an Italian cohort*. Allergy, 2015. 70(8): p. 933-43.
180. Schmid, J.M., et al., *Pretreatment IgE sensitization patterns determine the molecular profile of the IgG4 response during updosing of subcutaneous immunotherapy with timothy grass pollen extract*. J Allergy Clin Immunol, 2016. 137(2): p. 562-70.
181. Schmid-Grendelmeier, P., *[Recombinant allergens. For routine use or still only science?]*. Hautarzt, 2010. 61(11): p. 946-53.
182. Schuler, S., et al., *Microarray-based component-resolved diagnosis of latex allergy: isolated IgE-mediated sensitization to latexprofilin Hev b8 may act as confounder*. Clin Transl Allergy, 2013. 3(1): p. 11.
183. Schulten, V., et al., *Pru p 3, the nonspecific lipid transfer protein from peach, dominates the immune response to its homolog in hazelnut*. Allergy, 2011. 66(8): p. 1005-13.
184. Schwarz, A., et al., *IgG and IgG4 to 91 allergenic molecules in early childhood by route of exposure and current and future IgE sensitization: Results from the Multicentre Allergy Study birth cohort*. J Allergy Clin Immunol, 2016. 138(5): p. 1426-1433 e12.
185. Seyfarth, F., et al., *Diagnosing latex sensitizations with recombinant latex allergens - the Immunosolid-phase allergen chip (ISAC (R))*. 2011. 163-163.
186. Seyfarth, F., et al., *Diagnostic value of the ISAC(R) allergy chip in detecting latex sensitizations*. Int Arch Occup Environ Health, 2014. 87(7): p. 775-81.
187. Shamji, M.H., et al., *Functional rather than immunoreactive levels of IgG4 correlate closely with clinical response to grass pollen immunotherapy*. Allergy, 2012. 67(2): p. 217-26.
188. Shoda, T., et al., *Yogurt consumption in infancy is inversely associated with atopic dermatitis and food sensitization at 5 years of age: A hospital-based birth cohort study*. J Dermatol Sci, 2017. 86(2): p. 90-96.
189. Shreffler, W.G., *Microarrayed recombinant allergens for diagnostic testing*. J Allergy Clin Immunol, 2011. 127(4): p. 843-9; quiz 850-1.
190. Simon, D., et al., *Frequent sensitization to *Candida albicans* and profilins in adult eosinophilic esophagitis*. Allergy, 2013. 68(7): p. 945-8.

191. Simpson, A., et al., *Patterns of IgE responses to multiple allergen components and clinical symptoms at age 11 years*. J Allergy Clin Immunol, 2015. 136(5): p. 1224-31.
192. Siroux, V., et al., *Specific IgE and IgG measured by the MeDALL allergen-chip depend on allergen and route of exposure: The EGEA study*. J Allergy Clin Immunol, 2017. 139(2): p. 643-654 e6.
193. Skrindo, I., et al., *The use of the MeDALL-chip to assess IgE sensitization: a new diagnostic tool for allergic disease?* Pediatr Allergy Immunol, 2015. 26(3): p. 239-246.
194. Stemeseder, T., et al., *Cross-sectional study on allergic sensitization of Austrian adolescents using molecule-based IgE profiling*. Allergy, 2017. 72(5): p. 754-763.
195. Stemeseder, T., et al., *Exposure to Indoor Allergens in Different Residential Settings and Its Influence on IgE Sensitization in a Geographically Confined Austrian Cohort*. PLoS One, 2017. 12(1): p. e0168686.
196. Szepfalusi, Z., et al., *Preventive sublingual immunotherapy in preschool children: first evidence for safety and pro-tolerogenic effects*. Pediatr Allergy Immunol, 2014. 25(8): p. 788-95.
197. Thomas, W.R., *Hierarchy and molecular properties of house dust mite allergens*. Allergol Int, 2015. 64(4): p. 304-11.
198. TolKKI, L., et al., *Clinical characterization and IgE profiling of birch (*Betula verrucosa*)--allergic individuals suffering from allergic reactions to raw fruits and vegetables*. J Allergy Clin Immunol Pract, 2013. 1(6): p. 623-31 e1.
199. Tosca, M.A., et al., *Egg allergy: the relevance of molecular-based allergy diagnostics*. Clin Exp Allergy, 2014. 44(8): p. 1094-5.
200. Treudler, R., *Update on in vitro allergy diagnostics*. J Dtsch Dermatol Ges, 2012. 10(2): p. 89-97; quiz 98-9.
201. Treudler, R. and J.C. Simon, *Overview of component resolved diagnostics*. Curr Allergy Asthma Rep, 2013. 13(1): p. 110-7.
202. Tuppo, L., et al., *Pearmaclein-a new peach allergenic protein: similarities, differences and misleading features compared to Pru p 3*. Clin Exp Allergy, 2013. 43(1): p. 128-40.
203. Twaroch, T.E., et al., *Carrier-bound Alt a 1 peptides without allergenic activity for vaccination against *Alternaria alternata* allergy*. Clin Exp Allergy, 2012. 42(6): p. 966-75.
204. Ukleja-Sokolowska, N., et al., *Sunflower seed allergy*. Int J Immunopathol Pharmacol, 2016. 29(3): p. 498-503.
205. Van Gasse, A.L., et al., *Molecular allergy diagnosis: status anno 2015*. Clin Chim Acta, 2015. 444: p. 54-61.
206. van Hage, M., et al., *Performance evaluation of ImmunoCAP(R) ISAC 112: a multi-site study*. Clin Chem Lab Med, 2017. 55(4): p. 571-577.
207. van Kampen, V., et al., *IgE sensitization to lupine in bakers - cross-reactivity or co-sensitization to wheat flour?* Int Arch Allergy Immunol, 2015. 166(1): p. 63-70.
208. van Rhijn, B.D., et al., *Su1863 Component-Resolved Diagnostics Demonstrate Birch Pollen Sensitization With Cross-Reactivity to Food Allergens in Adult Eosinophilic Esophagitis*. Gastroenterology, 2013. 144(5): p. S-494-S-495.
209. van Rhijn, B.D., et al., *Birch pollen sensitization with cross-reactivity to food allergens predominates in adults with eosinophilic esophagitis*. Allergy, 2013. 68(11): p. 1475-81.
210. van Rhijn, B.D., et al., *Evaluation of allergen-microarray-guided dietary intervention as treatment of eosinophilic esophagitis*. J Allergy Clin Immunol, 2015. 136(4): p. 1095-7 e3.
211. Varga, E.M., et al., *Anaphylaxis to buckwheat in an atopic child: a risk factor for severe allergy to nuts and seeds?* Int Arch Allergy Immunol, 2011. 156(1): p. 112-6.
212. Verweij, M.M., et al., *Young infants with atopic dermatitis can display sensitization to Cor a 9, an 11S legumin-like seed-storage protein from hazelnut (*Corylus avellana*)*. Pediatr Allergy Immunol, 2011. 22(2): p. 196-201.
213. Verweij, M.M., et al., *Age-dependent sensitization to the 7S-vicilin-like protein Cor a 11 from hazelnut (*Corylus avellana*) in a birch-endemic region*. J Investig Allergol Clin Immunol, 2012. 22(4): p. 245-51.
214. Westman, M., et al., *Early childhood IgE reactivity to pathogenesis-related class 10 proteins predicts allergic rhinitis in adolescence*. J Allergy Clin Immunol, 2015. 135(5): p. 1199-206 e1-11.
215. Westwood, M., et al., *ImmunoCAP(R) ISAC and Microtest for multiplex allergen testing in people with difficult to manage allergic disease: a systematic review and cost analysis*. Health Technol Assess, 2016. 20(67): p. 1-178.
216. Vieira, T., et al., *Microarray based IgE detection in poly-sensitized allergic patients with suspected food allergy - an approach in four clinical cases*. Allergol Immunopathol (Madr), 2012. 40(3): p. 172-80.
217. Villalta, D., et al., *Isolated IgE reactivity to native walnut vicilin-like protein (nJug r 2) on ISAC microarray is due to cross-reactive carbohydrate epitopes*. Clin Chem Lab Med, 2013. 51(10): p. 1991-5.
218. Villalta, D., et al., *Allergen component specific ige measurement with the Immulite 2000 system: diagnostic accuracy and intermethod comparison*. J Clin Lab Anal, 2015. 29(2): p. 135-41.
219. Williams, P., et al., *Evaluation of a novel automated allergy microarray platform compared with three other allergy test methods*. Clin Exp Immunol, 2016. 184(1): p. 1-10.
220. Vitte, J., et al., *Multiplex FABER : application à l'allergie à la pêche, à propos de six cas méditerranéens*. Revue Française d'Allergologie, 2017. 57(3): p. 233-234.
221. Wohrl, S., et al., *The performance of a component-based allergen-microarray in clinical practice*. Allergy, 2006. 61(5): p. 633-9.
222. Wollmann, E., et al., *Natural clinical tolerance to peanut in African patients is caused by poor allergenic activity of peanut IgE*. Allergy, 2015. 70(6): p. 638-52.
223. Wollmann, E., et al., *Reduction in allergen-specific IgE binding as measured by microarray: A possible surrogate marker for effects of specific immunotherapy*. J Allergy Clin Immunol, 2015. 136(3): p. 806-809 e7.
224. Zeng, G., et al., *Component-Resolved Diagnostic Study of Dermatophagoides Pteronyssinus Major Allergen Molecules in a Southern Chinese Cohort*. J Investig Allergol Clin Immunol, 2015. 25(5): p. 343-51.
225. Zennaro, D., et al., *Proteomics plus genomics approaches in primary immunodeficiency: the case of immune dysregulation, polyendocrinopathy, enteropathy, X-linked (IPEX) syndrome*. Clin Exp Immunol, 2012. 167(1): p. 120-8.
226. Zheng, Y.W., et al., *Allergen micro-array detection of specific IgE-reactivity in Chinese allergy patients*. Chin Med J (Engl), 2011. 124(24): p. 4350-4.



Seit über 40 Jahren versorgen wir Ärzte in der ganzen Welt mit zuverlässigen Services – als Ihr Partner des Vertrauens, heute und in der Zukunft.

**Wünschen Sie weitere Informationen?**  
Sprechen Sie uns gerne an.

[thermoscientific.com/phadia/de](http://thermoscientific.com/phadia/de)

© 2017 Thermo Fisher Scientific Inc. Alle Rechte vorbehalten. Alle Warenzeichen sind das Eigentum von Thermo Fisher Scientific und seiner Tochtergesellschaften, falls nicht anders angegeben. Rechtmäßiger Hersteller: Phadia AB, Uppsala, Schweden

**Thermo Fisher Diagnostics GmbH**, Munzinger Str. 7, D-79111 Freiburg, Tel. +49 761 47 8050, Fax +49 761 47 805338

**Thermo Fisher Diagnostics Austria GmbH**, Dresdner Str. 89, A-1200 Wien, Tel. +43 1 270 20 20, Fax +43 1 270 20 20

**Thermo Fisher Diagnostics AG**, Sennweidstr. 46, CH-6312 Steinhausen, Tel. +41 43 343 40 50, Fax +41 43 343 40 51

12/2017

**ThermoFisher**  
SCIENTIFIC